

SHEVCHENKO SCIENTIFIC SOCIETY
COMMISSION OF MATHEMATICS

**MATHEMATICAL
BULLETIN**

of the SHEVCHENKO
SCIENTIFIC SOCIETY

Volume 8

Ivano-Frankivsk — 2011

НАУКОВЕ ТОВАРИСТВО ім. ШЕВЧЕНКА
МАТЕМАТИЧНА КОМІСІЯ

**МАТЕМАТИЧНИЙ
ВІСНИК**

НАУКОВОГО ТОВАРИСТВА
ім. ШЕВЧЕНКА

Том 8

Івано-Франківськ — 2011

НАУКОВЕ ТОВАРИСТВО ім. ШЕВЧЕНКА МАТЕМАТИЧНИЙ ВІСНИК

Головний редактор:
Editor in chief:

А.Самойленко (Київ)
A.Samoilenko (Kyiv)

Редактори: О.Артемович (Івано-Франківськ), А.Загороднюк (Івано-Франківськ), В.Маслюченко (Чернівці), І.Микитюк (Львів), М.Перестюк (Київ), І.Парасюк (Київ), А.Плічко (Кіровоград), А.Прикарпатський (Дрогобич), М.Притула (Львів), В.Самойленко (Київ), О.Станжицький (Київ), В.Шарко (Київ).

Editors: O.Artemovych (Ivano-Frankivsk), A.Zagorodnyuk (Ivano-Frankivsk), V.Maslyuchenko (Chernivtsi), I.Mykytyuk (Lviv), M.Perestyuk (Kyiv), I.Parasyuk (Kyiv), A.Plichko (Kirovograd), A.Prykarpatsky (Drohobych), M.Prytula (Lviv), V.Samoilenko (Kyiv), O.Stanzhuk (Kyiv), V.Sharko (Kyiv).

Редакційна колегія:

А.Августинович (Гданськ, Польща), Т.Банакх (Львів), Д.Блекмор (Нью-Джерсі, США), М.Блашак (Познань, Польща), М.Боголюбов(мол.) (Москва), Б.Винницький (Дрогобич), В.Гафійчук (Львів), П.Голод (Київ), М.Горбачук (Київ), Л.Горневич (Торунь, Польща), Р.Грогорчук (Москва), П.Гудивок (Ужгород), Ю.Дрозд (Київ), В.Євтухов (Одеса), Я.Єлейко (Львів), М.Зарічний (Львів), Ю.Зелінський (Київ), З.Камонт (Гданськ, Польща), В.Кириченко (Київ), М.Комарницький (Львів), Б.Копитко (Львів), О.Литвин (Харків), О.Лопушанський (Львів), Я.Микитюк (Львів), З.Перадзинський (Варшава, Польща), Р.Петришин (Чернівці), М.Працьовитий (Київ), Я.Прикарпатський (Київ), Б.Пташник (Львів), О.Скасків (Львів), Я.Слав'яновський (Варшава, Польща), М.Черніков (Київ).

Editorial board:

A.Augustynowicz (Gdańsk, Poland), T.Banakh (Lviv), D.Blackmore (NJ, USA), M.Blaszak (Poznań, Poland), M.Bogolyubov(jr.) (Moscow), B.Vynnytsky (Drohobych), V.Hafychuk (Lviv), P.Holod (Kyiv), M.Horbachuk (Kyiv), L.Górniewicz (Toruń, Poland), R.Hryhorchuk (Moscow), P.Hudyvok(Uzhgorod), Yu.Drozd (Kyiv), V.Evtukhov (Odesa), Ya.Yeleyko (Lviv), M.Zarichnyi (Lviv), Yu.Zelinsky (Kyiv), Z.Kamont (Gdańsk, Poland), V.Kyrychenko (Kyiv), M.Komarnytsky (Lviv), B.Kopytko (Lviv), O.Lopushansky (Lviv), O.Lytvyn (Kharkiv), Ya.Mykytyuk (Lviv), Z.Peradzynski (Warsaw, Poland), R.Petryshyn (Chernivtsi), M.Pratsiovyty (Kyiv), Ya.Prykarpatsky (Kyiv), B.Ptashnyk (Lviv), O.Skaskiv (Lviv), J.Slawianowski (Warszawa, Poland), M.Chernikov (Kyiv).

Відповідальні редактори: О.Артемович, А.Загороднюк
Відповідальний секретар: С.Шарин
Executive Editors: O.Artemovych, A.Zagorodnyuk
Executive Secretary: S.Sharyn

Адреса: вул. Винниченка 26, Львів 79008, Україна
Address: 26 Vynnychenko Str., Lviv 79008, Ukraine

Друкується за ухвалами Математичної комісії та Видавничої
ради Наукового товариства ім. Шевченка у Львові

ISSN 1812-6774

©Наукове товариство ім. Шевченка, 2011.

C O N T E N T S

Dmytro BODNAR, Maria BUBNYAK	
On convergence of 1-periodic branched continued fraction of special form	5
Taras VASYLYSHYN	
Construction of complex-valued functions by pre-defined values of integrals of their powers	17
Nataliya HOYENKO, Tamara ANTONOVA, Sergij RAKINTSEV	
Approximation for rations of Lauricella-Saran functions F_S with real parameters by branched continued fractions	28
Vasyl' DAVYDOVYCH	
Exact solutions of a reaction-diffusion equations system modeling the predator-prey interaction	43
Maryan DMYTRYSHYN	
Exponential type vectors of closed operators on tensor products	60
Roman DMYTRYSHYN	
About some regions of convergence of a multidimensional J -fraction with independent variables	69
Andriy ZAGORODNYUK, Zoryana MOZHYROVSKA	
Composition operators on a Hilbert space	78
Roman ZATORSKY	
Paradeterminant product of triangle matrices and its properties	87
Myhailo KOPACH, Anatolij OBSHTA, Bohdan SHUVAR	
The iterative aggregation for non-linear operator equations	99

Viktoria KRAVTSIV

Algebra of block-symmetric polynomials: generating elements and translation operator 107

Andriy LOPUSHANSKY

The regularity of abstract Cauchy problem solutions for semi-linear parabolic equation 122

Volodymyr MASLYUCHENKO, Vasyl' NESTERENKO

Decomposition of continuity and transition maps 132

Bogdana OLIYNYK

Metric realization of regular permutation groups 151

Mykhailo OSYPOCHUK, Mykola PORTENKO

A Limit distribution for local times at zero of a sequence of diffusion processes 167

Aleksandr SAVCHENKO

Functors on the category of K -ultrametric spaces 184

Andriy SEMECHUK

Estimation of convergence error of the rational estimations of recurrence single periodical fractions of 3-d order to their values ... 199

Oleh SKASKIV, Olga ZADOROZHNA

Inequalities of Viman type without exceptional sets for double entire Dirichlet series 207

Ivan TVERDOKHLIB, Myroslava VOVK, Mykola PRYTULA, Yarema PRYKARPATSKY

An economical equilibrium consumption mathematical model with fuzzy given parameters of an associated financial network 217

Sergii SHARYN

Shift semigroups in the algebra of polynomial Schwartz tempered
distributions 230

Roman SHEVCHUK

Inhomogeneous diffusion processes on a half-line, generated by the
differential operator with Feller-Wentzel boundary condition ... 243

Denis BLACKMORE, Yarema PPRYKARPATSKY, Jolanta GOLENIA,
Anatoliy PRYKARPATSKY

The AKNS hierarchy and the Gurevich-Zybin dynamical system
integrability revisited 258

Igor GAPYAK, Viktor GERASIMENKO

On the evolution of observables and the Enskog kinetic
equation 283

Oleh NYKYFORCHYN, Oksana MYKYTSEY

L -idempotent linear operators between predicate semimodules, dual
pairs and conjugate operators 299

Michael OLEKSIENKO

Hardy type spaces on reduced Heisenberg groups 315

Lilia SKASKIV, Orest ARTEMOVYCH

Groups with many Černikov-by-nilpotent subgroups 328

Oleh SKASKIV, Andriy KURYLIAK

The probability of absence zeros in the disc for some random analytic
functions 335

GROUPS WITH MANY ČERNIKOV-BY-NILPOTENT SUBGROUPS

©2011 *Lilia SKASKIV, Orest ARTEMOVYCH*

Vasyl Stefanyk Precarpathian National University,
57 Shevchenka Str., Ivano-Frankivsk 76018, Ukraine,
Institute of Mathematics, Cracow University of Technology,
24 Warszawska Str., Cracow 31155, Poland

e-mail: *orest_artemovych@hotmail.com, lila_yonyk@ua.fm*

Received 27 August 2011

We prove that a non-perfect group is a minimal non-“Černikov-by-nilpotent” group if and only if it is a Heineken-Mohamed type group.

Introduction

Let χ be a class of groups closed under subgroups. A minimal non- χ group G is a group which is not a χ -group, while all proper subgroups of G are χ . Belyaev and Sesekin (see [1] and [2]) have determined the minimal non-“finite-by-abelian” groups. Bruno and Phillips [3] have classified infinite groups in which every proper subgroup is finite-by-“nilpotent of class c ”. Later Otal and Peña [4] have extended this class of groups by replacing the term “finite group” with “Černikov group” and have considered the locally graded groups in which every proper subgroup is Černikov-by-nilpotent.

УДК: 519.41/47; MSC 2000: 20F16, 20F19

Key words and phrases: nilpotent group, Černikov group, Heineken-Mohamed type group

In particular, it was proved that a locally graded group G is Černikov-by-“nilpotent of class c ” if and only if all proper subgroups of G are Černikov-by-“nilpotent of class c ” ($c \geq 1$ is an integer). Xu [5] has investigated the minimal non-“finite-by-nilpotent” groups.

In this paper we study the groups with many Černikov-by-nilpotent subgroups. We concerned the non-perfect groups in which all proper subgroups are Černikov-by-nilpotent and prove the following

Theorem. *Let G be a non-perfect group. Then G is a minimal non-“Černikov-by-nilpotent” group if and only if it is a Heineken-Mohamed type group.*

Minimal non-nilpotent groups with subnormal proper subgroups (so-called the Heineken-Mohamed type groups) are constructed, for example, in [6].

Throughout this paper p is a prime, $\mathbb{Z}$ is the ring of integers. For a group G , we denote by $Z(G)$ the centre, by $G', G'', \dots, G^{(n)}, \dots$ the members of the derived series, by $\gamma_1(G) = G, \gamma_2(G), \dots, \gamma_m(G), \dots$ the members of the lower central series of G . Moreover $\mathbb{Z}G$ means the group ring of G over $\mathbb{Z}$ and $H \triangleleft G$ means that H is a normal subgroup in G . Recall that a group G is non-perfect if $G' \neq G$.

We shall also use other standard terminology from [7] and [8].

1 Minimal non-“Černikov-by-nilpotent” groups

From Theorem 1 of [4] it follows that a locally graded group G with Černikov-by-abelian proper subgroups is Černikov-by-abelian. Recall that a group G is called indecomposable if any two proper subgroups of G generate a proper subgroup in G , and is called decomposable otherwise.

In the sequel we shall need the next lemmas.

Lemma 1.1. *Let G be a non-perfect group. If all proper normal subgroups of G are Černikov-by-nilpotent, then one of the following statements holds:*

- (1) G is a Černikov-by-nilpotent group;
- (2) G/G' is a quasicyclic group and G' is a torsion subgroup;

(3) G/G' is a cyclic group.

Proof. If the quotient group G/G' is decomposable, then $G = AB$ is a product of two proper normal subgroups A and B . Since A and B are Černikov-by-nilpotent, G is such as well.

Now let G/G' be an indecomposable group. Then by Lemma 2 of [9] G/G' is a quasicyclic p -group or a cyclic p -group for some prime p .

Suppose that G is not a Černikov-by-nilpotent group and the quotient group G/G' is quasicyclic. We need to show that the derived subgroup G' is torsion. For this, assume by contrary that G' is non-torsion. Then without restricting of generality we can assume that G' is an abelian torsion-free subgroup.

Let q be a prime different from p . By Lemma 2.3 of [10] there exists a proper $\mathbb{Z}[G/G']$ -submodule N in G' such that $\overline{G'} = G'/N$ is a q -group. If a is any element of G and $\overline{G} = G/N$, then $\overline{G'}\langle\overline{a}\rangle$ is an abelian group, and this gives a contradiction. Hence G' is a torsion subgroup. The proof is completed. $\square$

Lemma 1.2. *Let G be a non-perfect group with Černikov-by-nilpotent proper subgroups. If G/G' is a cyclic p -group, then G is a Černikov-by-nilpotent group or the derived subgroup G' is torsion.*

Proof. Let G be a non-“Černikov-by-nilpotent” group. Suppose that the result is false. Then without loss of generality we may assume that G' is an abelian torsion-free subgroup. Obviously $G = G'\langle a \rangle$ for some element $a \in G$, where $a^{p^n} \in G'$ for some positive integer n . Let q and r be the distinct primes different from p . By Lemma 2.3 of [10] there exists a proper $\mathbb{Z}[G/G']$ -submodule M of G' such that

$$\overline{G'} = G'/M = \overline{A} \times \overline{B}$$

is a group direct product of a non-trivial q -subgroup $\overline{A}$ and a non-trivial r -subgroup $\overline{B}$. Let A (and respectively B) be an inverse image of $\overline{A}$ (and respectively $\overline{B}$) in G . Then $A\langle a \rangle$ and $B\langle a \rangle$ are two proper subgroups of G and therefore there are positive integers k and s such that $\gamma_k(A\langle a \rangle) \leq A$, $\gamma_s(B\langle a \rangle) \leq B$ and $\gamma_k(A\langle a \rangle)$, $\gamma_s(B\langle a \rangle)$ are Černikov. This gives that G is Černikov-by-nilpotent, which is a contradiction. $\square$

Lemma 1.3. *Let G be a group with the quasicyclic quotient group G/G' . Then G is a minimal non-“Černikov-by-nilpotent” group if and only if G is a Heineken-Mohamed type group.*

Proof. $(\Leftarrow)$ is clear.

$(\Rightarrow)$ It is now easy to see that G does not contain a proper subgroup of finite index. Let S be a proper subgroup of G , $\overline{G} = G/G''$ and $\overline{S} = SG''/G''$. If $\overline{S}$ is proper in $\overline{G}$, then $\overline{S}$ (and consequently S) is a nilpotent group. Assume that $\overline{S} = \overline{G}$. Then there exists a positive integer k such that $\gamma_k(\overline{G})$ is a Černikov group and so $\gamma_k(\overline{G}) \leq Z(\overline{G})$. This means that $\overline{G}$ is a nilpotent group, a contradiction. $\square$

Lemma 1.4. *If G is a Černikov-by-nilpotent group, then G is Černikov or the derived subgroup G' is of infinite index in G .*

Proof. By the hypothesis, there is a positive integer m such that $\gamma_m(G)$ is a Černikov subgroup. If $\overline{G} = G/\gamma_m(G)$ is a finite group, then G is Černikov. Suppose that $\overline{G}$ is an infinite group. Then the quotient group $\overline{G}/\overline{G}'$ is also infinite and this completes the proof. $\square$

Newman and Wiegold [11] have determined the structure of minimal non-nilpotent groups with maximal subgroups (see also Theorem 3.1 from [12]). By Theorem 2.5 of [12] every infinite soluble minimal non-nilpotent group is either of Heineken-Mohamed type or contains a maximal subgroup. Theorem 3.1 of [12] yields that a minimal non-nilpotent group with a maximal subgroup is a hypercentral Černikov-by-nilpotent group.

Lemma 1.5. *Let G be a group with the cyclic quotient p -group G/G' . If all proper subgroups of G are Černikov-by-nilpotent, then G is Černikov-by-nilpotent.*

Proof. Let G be as given. By hypothesis, $G = G'\langle a \rangle$ for some element $a \in G$, where $a^{p^n} \in G'$ for some positive integer n . Assume that G is not a Černikov-by-nilpotent group. Then in view of Lemma 1.2 the derived subgroup G' is a torsion subgroup.

Next we prove that G' does not contain a proper subgroup of a finite index. Assume to the contrary, i.e. let H be a proper G -invariant subgroup of finite index in G' . Then $H\langle a \rangle$ is a proper subgroup of G and so there

is a positive integer s such that $\gamma_s(H\langle a \rangle)$ is a Černikov subgroup. Since G' is not Černikov, we obtain in view of Lemma 1.4 that G'' is of infinite index in G' . Put $\overline{G} = G/G''$. Assume that $\overline{H}\langle \overline{a} \rangle = (H\langle a \rangle G'')/G''$ is a Černikov group, then $\overline{G}$ is a Černikov group as well. Inasmuch as $\gamma_m(G')$ is a Černikov subgroup for some positive integer m , $\gamma_m(G') \leq G''$ and $G'/\gamma_m(G')$ is a nilpotent group, we see in view of Corollary (see [13], p. 19) that $G'/\gamma_m(G')$ (and consequently G') is a Černikov group, a contradiction. By Lemma 1.4 this means that $(\overline{H}\langle \overline{a} \rangle)'$ is of infinite index in $\overline{H}\langle \overline{a} \rangle$. But then $\overline{G}/(\overline{H}\langle \overline{a} \rangle)'$ is a central-by-finite group, a contradiction with Theorem 4.12 of [8]. Hence G' is a $\mathcal{F}$ -perfect group. Moreover, $G'/\gamma_m(G')$ is a divisible abelian group by Theorem 2.2 of [7]. Hence $G'' = \gamma_m(G')$. By Theorem 3.29 of [8] $G' = C_{G'}(G'')$ and therefore G' is a divisible abelian group.

Let D be a quasicyclic subgroup of G' such that $D \not\subseteq Z(G)$ and $T = \langle D, a \rangle$. Assume that $T \neq G$. Then $\gamma_k(T)$ is a Černikov subgroup for some positive integer k and $\hat{T} = T/\gamma_k(T)$ is a nilpotent group. Inasmuch $\hat{D} \subseteq Z(\hat{T})$, we deduce that $\hat{T}$ is an abelian group and therefore $T' = \gamma_k(T)$. But then $T'\langle a \rangle$ is Černikov and so

$$D \leq C_T(T'\langle a \rangle).$$

This yields that T is an abelian group, a contradiction. Thus we conclude that $T = G$ and $\langle D^x \mid x \in G \rangle = G'$. Since $C_G(D)$ is of finite index in G , G' is a Černikov group. This final contradiction proves the lemma. $\square$

Proof of Theorem. Follows from Lemmas 1.1, 1.3 and 1.5. $\square$

- [1] Belyaev V.V., Seseikin N.F. On infinite groups of Miller-Moreno type // Acta Math. Acad. Sci. Hungar. — 1975. — **26**. — P. 369–376. (in Russian)
- [2] Belyaev V.V. Groups of Miller-Moreno type // Siberian Math. Zh. — 1978. — **19**. — P. 509–514. (in Russian)

- [3] *Bruno B., Phillips R.E.* On minimal conditions related to Miller-Moreno type groups // *Rend. Sem. Mat. Univ. Padova.* — 1983. — **69**. — P. 153-168.
- [4] *Otal J., Peña J.M.* Groups in which every proper subgroup is Černikov-by-nilpotent or nilpotent-by-Černikov // *Archiv Math.* — 1988. — **51**. — P. 193-197.
- [5] *Xu M.* Groups whose proper subgroups are finite-by-nilpotent // *Archiv Math.* — 1996. — **66**. — P. 353-359.
- [6] *Heineken H., Mohamed I.J.* A group with trivial centre satisfying the normalizer condition // *J. Algebra.* — 1968. — **10**. — P. 368-376.
- [7] *Černikov S.N.* Groups with prescribed properties of system of groups. — Moskow: Nauka, 1978. — 384 p. (in Russian)
- [8] *Robinson D.J.S.* Finiteness Conditions and Generalized Soluble Groups, Part 1. — Berlin Heidelberg New York: Springer-Verlag, 1972. — 210 p.
- [9] *Artemovych O.D.* On indecomposable metabelian groups // *Ukrainian Math. Zh.* — 1990. — **42**. — P. 1252-1254. (in Russian)
- [10] *Bruno B., Phillips R.E.* A note on groups with nilpotent-by-finite proper subgroups // *Archiv Math.* — 1995. — **65**. — P. 369-374.
- [11] *Newman M.F., Wiegold J.* Groups with many nilpotent subgroups // *Archiv Math.* — 1964. — **15**. — P. 241-250.
- [12] *Xu M.* Groups whose proper subgroups are Baer groups // *Acta Math. Sinica, New Series.* — 1996. — **12**. — P. 10-17.
- [13] *Robinson D.J.S.* Infinite soluble and nilpotent groups. — London: Queen Mary College Math. Notes, 1967. — 226 p.

**ГРУПИ, БАГАТІ
НІЛЬПОТЕНТНИМИ-НАД-ЧЕРНІКОВСЬКИМИ
ПІДГРУПАМИ**

Ліля СКАСКІВ, Орест АРТЕМОВИЧ

Прикарпатський національний університет імені Василя Стефаника,
вул. Шевченка 57, Івано-Франківськ 76018, Україна,
Краківська Політехніка,
вул. Варшавська 24, Краків 31155, Польща
e-mail: *orest_artemovych@hotmail.com, lila_yonuk@ua.fm*

Доведено, що недосконала група є мінімальною не нільпотентною-над-черніковською тоді і тільки тоді, коли вона група типу Хайнекена-Мохамеда.